


Atty. Docket No. 94090

21. (original) A method according to claim 20, wherein the reaction is performed according to the Mitsunobu reaction protocol, to yield an ester derivative of the 3β -hydroxy- 5β -H steroidal sapogenin.
22. (original) A method according to claim 20, wherein the activated derivative of the sapogenin is an organic sulphonated derivative.
23. (original) A method for the synthesis of smilagenin, comprising catalytic hydrogenation of diosgenone followed by reduction of the resulting 3-keto, 5β -H steroidal sapogenin using a hindered organoborane.
24. (original) A method for the synthesis of epismilagenin, comprising catalytic hydrogenation of diosgenone followed by reduction of the resulting 3-keto, 5β -H steroidal sapogenin using anorganoaluminohydride.
25. (currently amended) A method according to ~~any one of the preceding claims~~ claim 20, wherein a sapogenin initially formed is subsequently converted to a pro-drug form thereof or to another physiologically acceptable form thereof.

Respectfully submitted,

WELSH & KATZ, LTD.



Gerald T. Shekleton
Registration No. 27,466

Dated: April 11, 2005

Welsh & Katz, Ltd.

120 South Riverside Plaza, 22nd Floor

Chicago, Illinois 60606

Telephone: (312) 655-1500